

339: WILKINS-CHALGREN ANAEROBE BROTH

This recipe contains strain-specific modifications for *Oliverpabstia intestinalis* DSM 106162 *

Final pH: 6.8 - 7.0

Final volume: 1000 ml

dehydrated Wilkins-Chalgren medium (Oxoid CM0643)	33.00	g
<u>Sodium resazurin (0.1% w/v)</u>	<u>0.50</u>	<u>ml</u>
L-Cysteine HCl	0.30	g
Phenosaftranine	2.00	ml/l
Cystein	0.05	g/ml
DTT	0.02	g/ml
Distilled water	1000.00	ml

1. Dissolve 33.0 g/l dehydrated Wilkins-Chalgren medium (Oxoid CM0643) in distilled water and add 0.5 ml/l Na-resazurin solution (0.1% w/v), bring to the boil and cool to room temperature while sparging with 100% N₂ gas. Add L-cysteine-HCl x H₂O (0.3 g/l), dispense the medium under same gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclave. After autoclaving check pH and adjust to pH 6.8 - 7.0, if necessary.

2. Note: In case resazurin is not colourless addition of 10 - 20 mg sodium dithionite per liter (e.g. from 5% (w/v) solution, freshly prepared under N₂ and filter-sterilized) can be used to completely reduce the medium.

* Instead of Na-resazurin add 0,002 ml/ml Phenosafranin (Stock solution 1,25 mg/ml in H₂O). After heating add 0,010 ml/ml Cystein-DTT (DL-Dithiothreitol)-solution (0,05 g/ml Cystein, 0,02 g/ml DTT).