Microorganisms



402: ALICYCLOBACILLUS MEDIUM

This recipe contains strain-specific modifications for Alicyclobacillus sp. DSM 11983 *

Final volume: 1000 ml

Solution A	1000.00	ml
Solution B	1.00	ml
Solution C	500.00	ml
MnSO ₄	10.00	mg/l

Sterilize separately. For liquid medium combine solution A (with 1000.0 ml distilled water) and solution B. For solid medium combine solution A (with 500 ml distilled water), solution B and solution C. For strains of A. cycloheptanicus add 5 g/l of yeast extract instead of 2 g/l.

Solution A

$CaCl_2 \times 2 H_2O$	0.25	g
$MgSO_4 \times 7 H_2O$	0.50	g
$(NH_4)_2SO_4$	0.20	g
Yeast extract	2.00	g
Glucose	5.00	g
KH_2PO_4	3.00	g
Distilled water (for liquid medium)	1000.00	ml
Distilled water (for solid medium)	500.00	ml
Adjust pH to 4.0		
Solution B Trace element solution SL-6	1.00	ml
Solution C		
Agar	15.00	g
Distilled water	500.00	ml
Trace element solution SL-6 (from medium 2		
$ZnSO_4 \times 7 H_2O$	0.10	g
$MnCl_2 \times 4 H_2O$	0.03	g
H_3BO_3	0.30	g

^{*} Plus $MnSO_4$ (10 μ g/ml)

Microorganisms

402: ALICYCLOBACILLUS MEDIUM



$CoCl_2 \times 6 H_2O$	0.20	g
CuCl ₂ x 2 H ₂ O	0.01	g
NiCl ₂ x 6 H ₂ O	0.02	g
$Na_2MoO_4 \times 2 H_2O$	0.03	g
Distilled water	1000.00	ml