

402: ALICYCLOBACILLUS MEDIUM

This recipe contains strain-specific modifications for *Neobacillus fumarioli* DSM 18237 *

Final pH: * 5.5

Final volume: 1000 ml

Solution A	1000.00	ml
Solution B	1.00	ml
Solution C	500.00	ml

Sterilize separately. For liquid medium combine solution A (with 1000.0 ml distilled water) and solution B. For solid medium combine solution A (with 500 ml distilled water), solution B and solution C. For strains of *A. cycloheptanicus* add 5 g/l of yeast extract instead of 2 g/l.

* Adjust pH 5.5

Solution A

CaCl ₂ x 2 H ₂ O	0.25	g
MgSO ₄ x 7 H ₂ O	0.50	g
(NH ₄) ₂ SO ₄	0.20	g
Yeast extract	2.00	g
Glucose	5.00	g
KH ₂ PO ₄	3.00	g
Distilled water (for liquid medium)	1000.00	ml
Distilled water (for solid medium)	500.00	ml

Adjust pH to 4.0

Solution B

Trace element solution SL-6	1.00	ml
------------------------------------	------	----

Solution C

Agar	15.00	g
Distilled water	500.00	ml

Trace element solution SL-6 (from medium 27)

ZnSO ₄ x 7 H ₂ O	0.10	g
MnCl ₂ x 4 H ₂ O	0.03	g
H ₃ BO ₃	0.30	g
CoCl ₂ x 6 H ₂ O	0.20	g

402: ALICYCLOBACILLUS MEDIUM

$\text{CuCl}_2 \times 2 \text{ H}_2\text{O}$	0.01	g
$\text{NiCl}_2 \times 6 \text{ H}_2\text{O}$	0.02	g
$\text{Na}_2\text{MoO}_4 \times 2 \text{ H}_2\text{O}$	0.03	g
Distilled water	1000.00	ml